



HHS Public Access

Author manuscript

J Acad Nutr Diet. Author manuscript; available in PMC 2017 August 20.

Published in final edited form as:

J Acad Nutr Diet. 2013 August ; 113(8): 1062–1068. doi:10.1016/j.jand.2013.04.021.

Rural Disparities in the Distribution of Policies that Support Healthy Eating in US Secondary Schools

Marilyn S. Nanney, PhD, MPH, RD [associate professor],

Department of Family Medicine and Community Health, Program in Health Disparities Research

Cynthia S. Davey, MS [research associate], and

Clinical and Translational Science Institute, Biostatistical Design and Analysis Center

Martha Y. Kubik, PhD, RN [associate professor]

School of Nursing, all at the University of Minnesota, Minneapolis

Abstract

The distribution of food and nutrition policies and practices from 28 US states representing 6,732 secondary schools was evaluated using data from the 2008 School Health Profiles principal survey. School policies and practices evaluated were: availability of low-nutrient, energy-dense (LNED) snacks/drinks; use of healthy eating strategies; banning food marketing; availability of fruits and vegetables; and food package sizes. For each school, school-level demographic characteristics (percentage of students enrolled in free/reduced-price meals, minority enrollment, and geographic location) were also evaluated. Schools in small town/rural locations had significantly fewer policies that support healthy eating strategies and ban food marketing, and were less likely to serve fruits and vegetables at school celebrations, have fruits and vegetables available in vending or school stores, and limit serving-size packages. Schools serving the highest percentage of minority students consistently reported the same or better school food environments. However, schools serving the highest percentage of low-income students had varied results: vending and LNED vending policies were consistently better and fruit and vegetable availability-related policies were consistently worse. Disparities in the distribution of policies and practices that promote healthy school food environments seem most pronounced in small town/rural schools. The data also support the need for continued reinforcement and the potential for expansion of these efforts in urban and suburban areas and schools with highest minority enrollment.

Keywords

Rural disparities; School nutrition; Food policy

The child nutrition and wic reauthorization Act of 2004 (CNRA 2004), a major federal legislative milestone to address childhood obesity, included a mandate to school districts participating in the school meal program to establish and implement policies addressing

Address correspondence to: Marilyn S. Nanney PhD, MPH, RD, Department of Family Medicine and Community Health, Program in Health Disparities Research, University of Minnesota, 717 Delaware St SE, Minneapolis, MN 55414. msnanney@umn.edu.

STATEMENT OF POTENTIAL CONFLICT OF INTEREST

No potential conflict of interest was reported by the authors.

nutrition at school by the start of the 2006–2007 school year. Evaluating the effectiveness of these school nutrition policies is a nationwide priority.¹ Research suggests that school food environments, especially competitive foods, contribute excess energy to children's diets.^{2,3} Estimates show that low-nutrient, energy-dense (LNE) competitive foods provide 171 kcal/day to the diets of middle school students and 219 kcal/day to high school students.⁴ Although evidence of the impact on student body mass index remains mixed,⁵ a few studies suggest inverse associations between youth overweight and school food environments when mandated by states⁶ and whether strong district wellness policies⁷ and local school policies and practices are in place.^{8,9}

Monitoring the distribution of school food policies and practices and evaluating effectiveness through a health disparities lens is important.¹⁰ Minority and low-socioeconomic groups are disproportionately affected by excess weight at all ages.¹¹ Obesity rates among rural youth are as much as 50% higher compared with their urban counterparts.^{12–14} Although recent (2001–2008) evidence suggests a leveling off of obesity prevalence among white students, it also points to a worsening disparity among nonwhite students, especially for severe obesity.¹⁵ Descriptive evaluations considering school obesity prevention policies and disparities have identified differences by geographic region of the country (eg, southern states)¹⁶ and across schools by poverty status and geographic location (eg, schools in rural communities).^{17,18} Two evaluation studies report more favorable behavioral outcomes for minority students with stronger state mandates for competitive foods in schools.^{19,20} Evidence-based recommendations directed at competitive foods that aim to reverse the obesity trajectory include: (a) establishing nutrition standards for competitive foods available at school,²¹ (b) incorporating collaborative strategies to encourage healthy eating at school,²² and (c) banning advertising and marketing of LNE foods to youth in schools.^{21,23} The extent that the adoption of these recommendations is evenly distributed across US schools is not known. The goal of this study was to compare the distribution of food and nutrition-related policies and practices across US middle and junior/senior high schools (6th to 12th grade) by geographic location, minority enrollment, and free/reduced-price school meal enrollment after implementation of the 2004 CNRA in 2006. This inquiry is especially relevant as schools prepare to respond to the Healthy, Hunger-Free Kids Act of 2010, which focuses on competitive foods sold at school with a rolling implementation timeline beginning 2012–2020.²⁴

METHODS

Dependent Variables and Source

The dependent variables for this cross-sectional study were school-level food and nutrition-related policy and practice questions collected as part of the 2008 School Health Profiles Principal Survey (Profiles), a surveillance tool sponsored by the Centers for Disease Control and Prevention (CDC), which has good validity and reliability.²⁵ Profiles include a biennial survey of public school principals of middle and junior/senior high schools collected by state education and health agencies. School response rates for individual states for 2008 ranged from 70% to 93%.²⁶ Most states give the CDC permission to share their de-identified dataset. For this study, states were individually contacted and asked to share their school

identified data sets for the purpose of linking the school-level demographic data (geographic location, minority enrollment, free/reduced-priced meal enrollment).

Data Collection

An e-mail letter of invitation was sent to all 50 states' Profiles coordinators from the principal investigator explaining the study purpose and requesting access to their existing 2008 school-level policy data. A signed data use agreement detailing data confidentiality was also included. State Profiles administrators (ie, Department of Education, Department of Health) had varying comfort levels with data-sharing, ranging from accepting the terms of agreement, to requiring additional paperwork, to engaging in an iterative process with the project data manager that accomplished the data linking without divulging school identifying data. Total contacts (ie, emails, phone calls) to state agencies to share and transfer their states' Profiles data ranged from 1 to 35 with an average of 10. Fifty-six percent of states (n = 28) contacted agreed to share their datasets. A map of the participating states is available at the project website (<http://z.umn.edu/schoolnutrition>). Seven states did not respond to multiple requests to participate (14%); eight refused to participate (16%), citing change in leadership, concerns that sharing the identified datasets violated agreements with participating schools, or inability to locate data due to staff turnover and loss of positions. Seven states (14%) sent de-identified data, which could not be used.

State nonresponse bias was assessed using publicly available state-level policy and practice data from the nonparticipating states. The absolute differences between participating and nonparticipating states in the median weighted percentage of schools reporting each nutrition policy item were less than 5% for all but two items. The two items were "allowed students to purchase non-fried vegetables" (20% vs 28%) and "collected suggestions on food preferences" (45% vs 55%). Altogether, these results suggest similar nutrition policy implementation for participating and nonparticipating states.

Scale Development

Food and nutrition-related policy and practice items from the Profiles principal survey were identified and grouped to represent three domains: availability of LNEED snacks and drinks, engaging in healthy eating strategies, and marketing of LNEED snacks and drinks. Four other policy and practice items were evaluated individually.

Availability of LNEED snacks and drinks—Ten items were identified and included whether the following were available for purchase in vending machines or school stores (VMSS): chocolate candy, candy, salty snacks not low in fat, cookies, cakes, crackers not low in fat, ice cream not low in fat, 2% or whole milk, frozen water ices or slushes that do not contain juice, soda pop or fruit drinks that are not 100% juice, sport drinks, and caffeinated foods or beverages. Responses were coded as yes=1, no=0. The summated scale score represented the total number of LNEED snacks and drinks available for purchase. Cronbach's α for this scale was .80. Schools without VMSS were excluded from this analysis.

Healthy eating strategies—Five items were identified and included whether the following strategies were used: strategic pricing of healthy snacks and drinks lower in cost and/or LNEED snacks and drinks higher in cost, suggestions collected from students and families, calorie information provided to students/families, student taste-testing of new products, and student visits to the cafeteria for learning. Responses were coded as yes=1, no=0. The summated scale score represented the total number of implemented healthy eating strategies. Cronbach's α for this scale was .61.

Marketing of LNEED snacks and drinks—Four items were identified and included whether the school prohibits advertisements for candy, fast food, or soft drinks in (1) school building (yes/no), (2) on school grounds (yes/no), (3) on school buses (yes/no), (4) in school publications (yes/no). Responses were coded as yes=1, no=0. The summated scale score represented the total number of banned marketing practices. Cronbach's α for this scale was .89.

Other policy/practice items not included in the scales—Four additional items were examined separately: fruits and/or vegetables available at school celebrations (almost always or always, rarely, or never), any VMSS availability (yes or no), fruits and/or vegetables available for purchase from VMSS (yes or no), and limited package/serving size of items sold in VMSS (yes or no).

Independent Variables and Sources

The independent variables for this study were school-level demographic variables: geographic location (town/rural, urban, suburban), percentage minority enrollment (ie, defined as racial and ethnic minorities), and free/reduced-price school meal enrollment. The source of the independent variables was the National Center for Education Statistics Public Elementary/Secondary School Universe Survey (NCES), which is publicly available and updated annually.²⁷

Twelve NCES-defined geographic designations were combined into three locations for easiest interpretation: city (n=1,232 schools, 18.3%), suburban (n=1,467, 21.8%), and town/rural (n=4,033, 59.9%). The number of minority students for each school was calculated by subtracting the number of white non-Hispanic students enrolled from the total student enrollment and then dividing by the total student enrollment to calculate percentage minority enrollment. Similarly, the number of students enrolled in the free/reduced-price meal program was divided by the total student enrollment to calculate percent free/reduced-price meal enrollment.

These variables were categorized using approximate quartile cutpoints rounded to the nearest percentile divisible by five. The lack of significant differences in policy prevalence between the middle two quartiles resulted in the decision to combine these quartiles into one medium level category while preserving categories for the upper and lower approximate quartiles of schools. Minority percent enrollment data was available for 6,696 schools and was categorized as follows: <5% (low) (n=1,180 schools, 17.6%), 5% to <50% (medium) (n=3,802, 56.8%), and 50% or more (high) minority enrollment (n=1,714, 25.6%). Free/reduced-price meal enrollment data was available for 6,421 schools and categorized as

follows: <20% (low) (n=1,533 schools, 23.9%), 20% to <60% (medium) (n=3,501, 54.5%), and 60% or more (high) free/reduced-priced meal enrollment (n=1,387, 21.6%). The relationship between minority enrollment percentile and free/reduced-price percentile was examined to confirm that these two variables were not surrogates for each other. The correlation coefficient was 0.55, indicating that the two variables contribute unique school-level demographic information.

Analysis

Cronbach's α was used to assess internal reliability of summated scale variables. Correlation analysis was used to examine the linear relationship between minority enrollment percentile and free/reduced-price meal enrollment percentile. Multiple logistic regression models were used to estimate adjusted odds ratios for individual nutrition policy and practice items by location, minority enrollment category, and free/reduced-price meal enrollment category. Generalized linear models were used to evaluate location, minority enrollment category, and free/reduced-price meal enrollment category differences in adjusted mean scores for the nutrition policy and practice scales. Significant factors in each model were further evaluated to identify levels of school characteristics with significant differences. The Bonferroni method was used to adjust significance levels for these multiple comparisons (α of .017 for means, α of .008 for odds ratios). Model estimates and standard errors were used to construct 95% CIs for the adjusted means for each level of the school characteristic and adjusted odds ratios for school characteristic levels relative to the reference level.

RESULTS AND DISCUSSION

A total of 6,732 schools from 28 states were included in the analysis. A table detailing the distribution of each policy and practice item by each demographic characteristic is available from the project website (<http://z.umn.edu/schoolnutrition>). Table 1 identifies the mean score (95% CI) for each scale by school location and demographic category adjusted for the other school characteristic variables. For the *Availability of LNE D Snacks and Drinks Policy Scale*, a lower score was the better result. The adjusted mean number of LNE D snacks or drinks available for purchase from VMSS was significantly less (ie, better) for schools with the highest free/reduced-price meal enrollment than for schools with low or medium enrollment and was also significantly less for schools with medium enrollment than for schools with low enrollment. Studies show that restricting the availability of LNE D foods in schools while increasing the availability of healthful foods might be an effective strategy for promoting more healthful choices among students at school.^{4,28}

For the *Healthy Eating Strategies Policy Scale*, a higher score was the better result. The adjusted mean number of healthy eating strategies implemented was significantly higher (ie, better) in urban and suburban schools than in town/rural schools. This finding is important because studies show that youth are sensitive to food pricing²⁹ and respond positively to taste testing healthy foods.³⁰

For the *Marketing of LNE D Snacks and Drinks Policy Scale*, a higher score was the better result. The adjusted mean number of locations in which advertisement for candy, fast food, or soft drinks was banned was significantly higher (ie, better) in urban and suburban schools

than in town/rural schools; and significantly higher in schools with high minority enrollment than in schools with low or medium minority enrollment. An increasing number of studies demonstrate direct causal effects of exposure to food advertising on young people's weight and higher rates of obesity.³¹

Table 2 identifies the individual policy and practice items by school location and demographic category adjusted for the other school characteristic variables. The likelihood of fruits and/or vegetables almost always or always being available at school celebrations was significantly higher at suburban schools than at town/rural schools, significantly lower at schools with low and medium minority enrollment than at schools with highest minority enrollment, and significantly higher at schools with lowest free/reduced-price meal enrollment than at schools with highest free/reduced-price meal enrollment.

The likelihood of having no vending or school store availability of foods was significantly lower in schools with low and medium minority enrollment than in schools with the highest minority enrollment, and was significantly lower at schools with low and medium free/reduced-price meal enrollment than at schools with the highest free/reduced-price meal enrollment.

The likelihood of fruits and/or vegetables being available for purchase from vending machines or school stores was significantly higher in urban and suburban schools than in town/rural schools, was significantly lower at schools with low and medium minority enrollment than at schools with highest minority enrollment, and was significantly higher at schools with low and medium free/reduced-price meal enrollment than at schools with highest free/reduced-price meal enrollment.

The likelihood of limited package or serving sizes of foods sold in school stores or vending was significantly higher in urban and suburban schools than in town/rural schools. Studies have identified an impact upon student dietary intake when portion sizes were limited in schools.^{32,33}

Strengths and Limitations

To our knowledge, this is the first study to compare a multistate sample of food and nutrition-related policies and practices across categories of place, ethnicity, and socioeconomic status after the implementation of the 2004 CNRA in 2006. The state response rate is a limitation. However, the authors made a reasonable attempt to determine whether bias exists. The school food and nutrition-related policies and practices evaluated were limited to those previously collected by states. No new data were collected.

The most pronounced disparity in the distribution of policies and practices that support healthy school food environments seems to exist among schools located in town and rural communities. These findings are troubling given the health and weight disparities that already exist among these communities. Living in rural areas is a risk factor for children being overweight or obese.¹² Explanations for this geographic difference supported by the literature include: smaller school size and therefore fewer resources, less availability of healthy foods, and higher cost of high-quality produce. A study of 14 rural Kansas high

schools reported that smaller food-service programs had fewer financial resources, fewer total products, and lower product volume, resulting in fewer fruit and vegetable options.¹³ Several studies have found that the availability and cost of healthy foods, especially high-quality produce, are most problematic in small town and rural settings.^{34–36}

Schools serving the highest percentage of minority students consistently reported the same or better school food environments. However, schools serving the highest percentage of low-income students had varied results; vending and LNEED vending policies were consistently better and fruit and vegetable availability related policies were consistently worse. A national multistate, multischool study investigating the school food environment reported better food and nutrition-related policies and practices among both lower income and higher minority enrollment schools.³⁷ However, this investigation occurred before the 2006 implementation of the 2004 CNRA. Another contradictory study involving a state examination of wellness policy language identified stronger policy language in the schools with the highest percentage of free/reduced-price meal enrollment.¹⁷ One plausible explanation for the counterintuitive findings is that schools serving high-needs areas have been targeted for supportive programming: coordinated school health, fresh fruit and vegetable programs, and summer foodservice programs.^{25,38,39} At the time of the current data collection, however, school-based fresh fruit and vegetable programs primarily targeted elementary schools and were in the piloting phase with limited reach. Continued monitoring of food policies and practices in secondary schools for geographic, income, and racial disparities is justified.

CONCLUSIONS

Students attending schools in small town and rural areas have significantly less exposure to healthy eating policies and practices and significantly more exposure to LNEED marketing at school than students attending urban and suburban schools. Students attending city and suburban schools and schools with the highest minority enrollment seem to be attending schools with better food environments, although the overall scale scores were low, indicating room for improvement. Findings from this study uniquely add to the literature in two ways. First, the evaluation period captures 2 years after the implementation of the mandatory school district wellness policies. Second, the assessments of policies and practices include those with a substantial evidence base not previously examined (eg, food marketing, promotional strategies). This post-CNRA implementation study strengthens the need for small town and rural focused policy supports, especially as next steps (the Healthy, Hunger-Free Kids Act of 2010) are implemented. The data also support the need for continued reinforcement and the potential for expansion of these efforts in urban and suburban areas and schools with the highest minority enrollment. Lastly, future research should examine the impact of these school food policy environments on student diet, weight, health, and academic outcomes. Especially important are evaluations by student sex, race/ethnicity, and income categories.

Acknowledgments

FUNDING/SUPPORT

Funding for this study was provided by the Robert Wood Johnson Foundation Healthy Eating Research Round 2 New Connections grant no. 65056.

Nancy Brener, PhD, Centers for Disease Control and Prevention, Division of Adolescent and School Health, provided substantial assistance with recruitment. Susan Lowry, Biostatistical Design and Analysis Center, Clinical and Translational Science Institute, performed the extensive data management activities. Deb Loy, RN, Minnesota Department of Education, provided insight into school food practices.

References

1. Story M, Nanney MS, Schwartz MB. Schools and obesity prevention: Creating school environments and policies to promote healthy eating and physical activity. *Milbank Q.* 2009; 87(1):71–100. [PubMed: 19298416]
2. Briefel RR, Crepinsek MK, Cabili C, Wilson A, Gleason PM. School food environments and practices affect dietary behaviors of US public school children. *J Am Diet Assoc.* 2009; 109(2 suppl):S91–S107. [PubMed: 19166677]
3. Kubik MY, Lytle LA, Hannan PJ, Perry CL, Story M. The association of the school food environment with dietary behaviors of young adolescents. *J Am Diet Assoc.* 2003; 93(7):1168–1173.
4. Fox MK, Gordon A, Nogales R, Wilson A. Availability and consumption of competitive foods in US public schools. *J Am Diet Assoc.* 2009; 109(2 Suppl):S57–S66. [PubMed: 19166673]
5. Jaime PC, Lock K. Do school based food and nutrition policies improve diet and reduce obesity? *Prev Med.* 2009; 48(1):45–53. [PubMed: 19026676]
6. Taber DR, Chriqui JF, Perna FM, Powell LM, Chaloupka FJ. Weight status among adolescents in states that govern competitive food nutrition content. *Pediatrics.* 2012; 130(3):437–444. [PubMed: 22891223]
7. Coffield JE, Metos JM, Utz RL, Waitzman NJ. A multivariate analysis of federally mandated school wellness policies on adolescent obesity. *J Adolesc Health.* 2011; 49(4):363–370. [PubMed: 21939866]
8. Fox MK, Dodd AH, Wilson A, Gleason PM. Association between school food environment and practices and body mass index of US public school children. *J Am Diet Assoc.* 2009; 109(2 suppl):S108–S117. [PubMed: 19166665]
9. Kubik MY, Lytle LA, Story M. Schoolwide food practices are associated with body mass index in middle school students. *Arch Pediatr Adolesc Med.* 2005; 159:1111–1114. [PubMed: 16330732]
10. Nanney M, Davey C. Evaluating the distribution of school wellness policies and practices: A framework to capture equity among schools serving the most weight vulnerable children. *J Am Diet Assoc.* 2009; 108(91):1436–1439.
11. Wang YF, Liang HF, Chen XL. Measured body mass index, body weight perception, dissatisfaction and control practices in urban, low-income African American adolescents. *BMC Public Health.* 2009; 9:183. [PubMed: 19523206]
12. Lutfiyya MN, Lipsky MS, Wisdom-Behounek J, Inpanbutr-Martinkus M. Is rural residency a risk factor for overweight and obesity for U.S. children? *Obesity.* 2007; 15(9):2348–2356. [PubMed: 17890504]
13. Nollen N, Bafort C, Davis A, et al. Competitive foods in schools: Availability and purchasing in predominately rural small and large high schools. *J Am Diet Assoc.* 2009; 109(5):857–864. [PubMed: 19394472]
14. Tae-Seale, T., Chandler, C. Rural Healthy People 2010: A companion document to Healthy People 2010. Vol. 2. College Station, TX: The Texas A&M University System Health Science Center, School of Rural Public Health; 2003. Nutrition and overweight concerns in rural areas: A literature review. <http://srph.tamhsc.edu/centers/rhp2010/09Volume2nutrition.pdf> [Accessed April 15, 2013]
15. Madsen KA, Weedn AE, Crawford PB. Disparities in peaks, plateaus, and declines in prevalence of high BMI among adolescents. *Pediatrics.* 2010; 126:434–442. [PubMed: 20713482]
16. Taber DR, Chriqui JF, Chaloupka FJ. Geographic disparities in state and district policies targeting youth obesity. *Am J Prev Med.* 2011; 41(4):407–414. [PubMed: 21961468]
17. Metos J, Nanney MS. The strength of school wellness policies: One state's experience. *J Sch Health.* 2007; 77(7):367–372. [PubMed: 17680895]

18. Nanney MS, Bohner C, Friedrichs M. Poverty-related factors associated with obesity prevention policies in Utah secondary schools. *J Am Diet Assoc.* 2008; 108(7):1210–1215. [PubMed: 18589031]
19. Taber DR, Stevens J, Evenson KR, et al. State policies targeting junk food in schools: Racial/ethnic differences in the effect of policy change on soda consumption. *Am J Public Health.* 2011; 101(9): 1769–1775. [PubMed: 21778484]
20. Taber DR, Chriqui JF, Chaloupka FJ. Differences in nutrient intake associated with state laws regarding fat, sugar, and caloric content of competitive foods. *Arch Pediatr Adolesc Med.* 2012; 166(5):452–458. [PubMed: 22566546]
21. Pilant VB. Position of the American Dietetic Association: Local support for nutrition integrity in schools. *J Am Diet Assoc.* 2006; 106(1):122–133. [PubMed: 16390677]
22. School Nutrition Association. [Accessed April 15, 2013] Local Wellness Policy Recommendations. http://www.schoolnutrition.org/uploadedFiles_old/SchoolNutrition.org/Child_Nutrition/Local_School_Wellness_Policies/SNALocalWellnessPolicyGuidelinesFinal.pdf
23. Institute of Medicine. Committee on Food Marketing and the Diets of Children and Youth. In: McGinnis, JM, Gootman, JA., Kraak, VI., editors. *Food marketing to children and youth: Threat or opportunity?*. Washington, DC: National Academies Press; 2006.
24. [Accessed March 10, 2013] Healthy Hunger Free Kids Act of 2010 (S. 3307 [111th]). Dec 13, 2010 <http://www.gpo.gov/fdsys/pkg/BILLS-111s3307enr/pdf/BILLS-111s3307enr.pdf>
25. Centers for Disease Control and Prevention. [Accessed March 10, 2013] School Health Profiles. <http://www.cdc.gov/healthyyouth/profiles/index.htm>
26. Brener, ND., McManus, T., Foti, K., et al. [Accessed March 10, 2013] School Health Profiles 2008: Characteristics of Health Programs Among Secondary Schools. http://www.cdc.gov/healthyyouth/profiles/2008/profiles_report.pdf
27. National Center for Educational Statistics. [Accessed March 10, 2013] Common Core of Data Public Elementary/ Secondary School Universe Survey: School Year 2006–07. <http://nces.ed.gov/ccd/pubschuniv.asp>
28. Cullen KW, Thompson DI. Texas school food policy changes related to middle school a la carte/ snack bar foods: Potential savings in kilocalories. *J Am Diet Assoc.* 2005; 105(12):1952–1954. [PubMed: 16321604]
29. French SA, Story MT, Fuller C, Gerlach AF. Food environment in secondary schools: A la carte, vending machines, and food policies and practices. *Am J Public Health.* 2003; 93:1161–1167. [PubMed: 12835203]
30. Nanney MS, Olaleye TM, Wang Q. Incorporating a healthy reimbursable snack in an afterschool homework program for middle school students: A case study. *Health Ed Behav.* 2012; 39(2):127–130.
31. Chou SY, Rashad I, Grossman M. Fast-food restaurant advertising on television and its influence on childhood obesity. *J Law Econ.* 2008; 51(4):599–618.
32. Hartstein J, Cullen KW, Reynolds KD, et al. Impact of portion-size control for school a la carte items: Changes in kilocalories and macronutrients purchased by middle school students. *J Am Diet Assoc.* 2008; 108(1):140–144. [PubMed: 18156001]
33. Cullen KW, Watson K, Baranowski T, Baranowski JH, Zakeri I. Squire’s Quest: Intervention changes occurred at lunch and snack meals. *Appetite.* 2005; 45(2):148–151. [PubMed: 15927302]
34. Dunn RA, Sharkey JR, Lotade-Manje J, Bouhlal Y, Nayga RM Jr. Socio-economic status, racial composition and the affordability of fresh fruits and vegetables in neighborhoods of a large rural region in Texas. *Nutr J.* 2011; 10(6):1–10. [PubMed: 21208446]
35. Larson NI, Story MT, Nelson MC. Neighborhood environments: Disparities in access to healthy foods in the U.S. *Am J Prev Med.* 2009; 36(1):74–78. [PubMed: 18977112]
36. Liese AD, Weis KE, Pluto D, Smith E, Lawson A. Food store types, availability, and cost of foods in a rural environment. *J Am Diet Assoc.* 2007; 107(11):1916–1923. [PubMed: 17964311]
37. Finkelstein DM, Hill EL, Whitaker RC. School food environments and policies in US public schools. *Pediatrics.* 2008; 122(1):e251–e259. [PubMed: 18595970]
38. USDA Fruit and Vegetable Program. [Accessed March 10, 2013] <http://www.ers.usda.gov/topics/food-nutrition-assistance/child-nutrition-programs/usda-fruit-and-vegetable-program.aspx>

39. USDA Summer Food Service Program. [Accessed March 10, 2013] <http://www.fns.usda.gov/summer-food-service-program-sfsp>

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1 Adjusted mean scale scores of policies and practices by school demographic characteristics for a sample of US secondary schools^a

Policy Domains	Sample Mean±SD ^b	Sample (Min, Max)	Location			Minority Enrollment			Free/Reduced-Price Meal Enrollment		
			Urban	Suburban	Town/Rural	Low (<5%)	Medium (5% to <50%)	High (>50%)	Low (<20%)	Medium (20% to <60%)	High (>60%)
Availability of LINED^{c,d} Snacks and Drinks Lower score better	4.42±2.82	(0, 10)	4.28 (4.06, 4.51)	4.39 (4.19, 4.59)	4.20 (4.06, 4.34)	4.26 (4.01, 4.51)	4.48 (4.33, 4.63)	4.13 (3.89, 4.38)	5.02 ^{**} (4.76, 5.28)	4.29 ^{**} (4.15, 4.43)	3.57 (3.33, 3.81)
Healthy Eating Strategies^c Higher score better	1.42±1.32	(0, 5)	1.53 (1.44, 1.61)	1.51 (1.43, 1.58)	1.36 ^{**} (1.31, 1.41)	1.46 (1.37, 1.55)	1.52 (1.47, 1.57)	1.41 (1.34, 1.48)	1.54 (1.46, 1.61)	1.44 (1.39, 1.49)	1.42 (1.33, 1.50)
Marketing of LINED Snacks and Drinks^c Higher score better	2.60±1.64	(0, 4)	2.75 (2.65, 2.85)	2.79 (2.70, 2.88)	2.48 ^{**} (2.42, 2.54)	2.47 ^{**} (2.36, 2.58)	2.69 [*] (2.63, 2.76)	2.85 (2.77, 2.94)	2.71 (2.61, 2.80)	2.60 (2.53, 2.67)	2.71 (2.61, 2.82)

^aN=2008 School Health Profiles data representing 6,732 schools from 28 states.

^bSD=standard deviation.

^cMean scores (95% confidence intervals), adjusted for other school-level characteristics, are reported for each scale.

^dLINED=low-nutrient, energy-dense. Includes chocolate candy, candy, salty snacks not low in fat, ice-cream not low in fat, cookies, cakes, crackers not low in fat, 2% or whole milk, water ices that do not contain juice, soda pop or fruit drinks that are not 100% juice, sport drinks, and caffeinated foods or beverages available for purchase in school vending machines or stores. The adjusted mean scores for the availability of LINED snacks and drinks scale are also adjusted for a significant interaction between minority enrollment and free/reduced-price meal eligibility.

^{*} Significantly worse scores than one or more location, minority enrollment, or free/reduced-price meal enrollment categories, adjusted for multiple comparisons from 0 at the 0.01 level.

^{**} Significantly worse scores than one or more location, minority enrollment, or free/reduced-price meal enrollment categories, adjusted for multiple comparisons from 0 at the 0.001 level.

Table 2
Adjusted odds ratio^a for each policy and practice by demographic characteristic for a sample of US secondary schools^b

Individual Policy items	Location			Minority Enrollment			Free/Reduced-Price Meal Enrollment		
	Urban	Suburban	Town/ Rural	Low (<5%)	Medium (5% to <50%)	High (>50%)	Low (<20%)	Medium (20% to <60%)	High (>60%)
Fruits and/or vegetables available at school celebrations ^c	1.17 (1.003, 1.36)	1.24* (1.08, 1.42)	ref ^e	0.60* (0.49, 0.72)	0.67* (0.58, 0.78)	ref	1.32* (1.10, 1.59)	1.14 (0.98, 1.32)	ref
No VMSS ^d availability of foods	1.09 (0.93, 1.28)	0.94 (0.81, 1.10)	ref	0.66* (0.54, 0.81)	0.64* (0.55, 0.75)	ref	0.60* (0.49, 0.72)	0.62* (0.54, 0.73)	ref
Fruits and/or vegetables are available for purchase from VMSS	1.48* (1.25, 1.76)	1.67* (1.43, 1.95)	ref	0.70* (0.57, 0.88)	0.78* (0.66, 0.93)	ref	2.26* (1.83, 2.79)	1.52* (1.27, 1.82)	ref
Limited package serving size of items sold in VMSS ^f	1.43* (1.19, 1.72)	1.64* (1.39, 1.93)	ref	0.79 (0.63, 0.98)	0.87 (0.72, 1.04)	ref	0.95 (0.77, 1.18)	1.03 (0.86, 1.23)	ref

^aOdds ratio and (95% confidence intervals), adjusted for other school-level characteristics for each nutrition policy item.

^bN=2008 School Health Profiles data representing 6,732 schools from 28 states.

^cIncludes responses “almost always or always”.

^dVMSS=vending machines or school stores.

^eref=reference group.

^fFor example: Does this school limit the package or serving size of any individual food and beverage items sold in vending machines or at the school store?

* Odds ratios significantly different from the reference group after adjusting for multiple comparisons at the 0.008 level.