

Location(s) of standing order use: MRI Departments of UAB Hospital & Ambulatory Clinics

MRI Standing Orders:

Hyoscyamine Administration for Use as Diagnostic Aid for Pelvic MRIs

Implementation Criteria: Patient scheduled for pelvic MRI or pelvic PET/MRI arrives for testing in MRI

Department. Personnel Authorized to Implement: MRI or PET/MRI Technologist or Registered Nurse in the Department of Radiology Orders:			
			Verify patient per the Patient Identification policy.
		2.	Verify provider's orders for pelvic MRI or PET/MRI and explain procedure to patient.
3.	Verify patient's allergies. If no allergy to hyoscyamine exist, administer hyoscyamine 0.25 mg SL (2 tablets of 0.125 mg dosage) immediately prior to MRI. If an allergy to hyoscyamine exists, do not administer hyoscyamine.		
4.	Evaluate patient for any side effects prior to release following testing.		
5.	Contact abdominal MRI attending if any side effects/patient complaints.		

Author: Ashley Thomas

Medical Director: Mason Frazier, MD 10/08/24

Vice Chair of Quality and Patient Experience: Samuel Galgano, MD

7/29/24

Pharmacy & Therapeutics Endorsement: 10/02/24 CNO Endorsement: Terri Poe DNP, RN 10/14/24 Medical Executive Committee Endorsement: 10/22/24 SO 264

Cross Reference: Generation of Orders Policy